

A global $SU(3)/U(3)$ flavor symmetry analysis for $B \rightarrow PP$ with $\eta - \eta'$ Mixing

Yu-Kuo Hsiao^{1,2,3}, Chia-Feng Chang⁴ and Xiao-Gang He^{5,4,2}

¹*Chongqing University of Post & Telecommunications, Chongqing, 400065, China*

²*Physics Division, National Center for Theoretical Sciences, Hsinchu, Taiwan 300*

³*Department of Physics, National Tsing Hua University, Hsinchu, Taiwan 300*

⁴*Department of Physics, National Taiwan University, Taipei, Taiwan 107*

⁵*Department of Physics and Astronomy,
Shanghai Jiao Tong University, Shanghai, China*

Abstract

A large number of new experimental data on B decay into two light pseudoscalar (P) mesons have been collected by the LHCb collaboration. Besides confirming information on $B_{u,d} \rightarrow PP$ decays obtained earlier by B-factories at KEK and SLAC, new information on $B_s \rightarrow PP$ and also more decay modes with P being η or η' have been obtained. Using these new data, we perform a global fit for $B \rightarrow PP$ to determine decay amplitudes in the framework of $SU(3)/U(3)$ flavor symmetry. We find that $SU(3)$ flavor symmetry can explain data well. The annihilation amplitudes are found to be small as expected. Several CP violating relations predicted by $SU(3)$ flavor symmetry are in good agreement with data. Current available data can give constraints on the amplitudes which induce $P = \eta, \eta'$ decays in the framework of $U(3)$ flavor symmetry, and can also determine the $\eta - \eta'$ mixing angle θ with $\theta = (-18.4 \pm 1.2)^\circ$. Several $B \rightarrow PP$ decay modes which have not been measured are predicted with branching ratios accessible at the LHCb. These decays can provide further tests for the framework of $SU(3)/U(3)$ flavor symmetry for B decays.

I. INTRODUCTION

A large number of experimental data on B decay into two pseudoscalar (P) mesons have been collected by the LHCb collaboration. Besides confirming information on $B_{u,d} \rightarrow PP$ obtained earlier by B-factories at KEK and SLAC, new information on B_s decays have been obtained [1, 2]. The new information can provide more insight about interactions responsible for B decays. $B \rightarrow PP$ are induced at one loop level in the standard model (SM). These decay modes being rare ones are expected to be sensitive to new physics beyond the SM. Before claiming the existence of any new physics beyond it is necessary to have the SM interactions be well understood. $B \rightarrow PP$ decays have been studied extensively in different ways. The main methods are QCD based perturbative calculations [3–5] and $SU(3)$ flavor symmetry [6–14].

The $SU(3)$ flavor symmetry approach has the advantage of being detailed dynamics independent. The decays are described by several $SU(3)$ invariant amplitudes which can lead to relations between different decay modes, but this approach by it-self cannot determine the size of the amplitudes. The QCD based perturbative approach being dynamic models, for example, the QCD factorization (QCDF) [3], perturbative QCD (pQCD) [4], and soft-collinear effective theory (SCET) [5], can calculate the very precisely measured CP violation asymmetry $\mathcal{A}_{CP}(\bar{B}^0 \rightarrow \pi^+ K^-) = (-8.2 \pm 0.6)\%$ [1, 2] for $\bar{B}^0 \rightarrow \pi^+ K^-$ decay. If the theory is universally valid they should be able to make accurate predictions for CP violation in other $B \rightarrow PP$ decays. These methods, however, all predict $\mathcal{A}_{CP}(\bar{B}^0 \rightarrow \pi^+ K^-) \approx \mathcal{A}_{CP}(B^- \rightarrow \pi^0 K^-)$, which is in contradiction with experimental observation. Therefore $\mathcal{A}_{CP}(\bar{B}^0 \rightarrow \pi^+ K^-) \neq \mathcal{A}_{CP}(B^- \rightarrow \pi^0 K^-)$ challenges these theories [15–17]. On the other hand, the analysis based on the $SU(3)$ flavor symmetry can be advantageous, where the different decay modes can be related and the relevant decay amplitudes be extracted from the data, despite of their unclear sources. A consistent solution for these CP violating asymmetries can be found. When sufficient data become available, the $SU(3)$ invariant amplitudes can be fixed and predictions be made, and the theory be tested. $SU(3)$ analysis may play a role to bridge dynamic theory and experimental data to provide some understanding of SM predictions for B decays.

The $SU(3)$ flavor symmetry has been wildly used for the studies in the SM for two-body and three-body mesonic B decays [18, 19], the extraction of the weak phase [20, 21], and the

constraint on new physics [22]. In its extended version, the two-body anti-triplet b -baryon decays of $\mathcal{B}_b \rightarrow \mathcal{B}_n M$ and $\mathcal{B}_b \rightarrow \mathcal{P}_c M$ decays can be studied [23–25], where \mathcal{B}_n and \mathcal{P}_c stand for the baryon and pentaquark state, respectively, with M the recoiled meson. To make sure $SU(3)$ flavor symmetry framework is valid for B decays, experimental test should be performed. Due to the fact that the Belle and Babar detectors at B-factories can only study B_u and B_d , but not B_s decays, the $SU(3)$ flavor symmetry have not been well tested. With the running of LHC, the LHCb has been able to obtain valuable data not only on $B_{u,d}$, but also B_s decays, one can therefore test more thoroughly the $SU(3)$ flavor symmetry for $B \rightarrow PP$ decays [26]. When more b -baryon decays are measured, $SU(3)$ can also be tested for the b -baryon sector. Experimentally, the data collections for the $B \rightarrow PP$ decays are in fact still not satisfactory. For example, $\bar{B}_s^0 \rightarrow K^0 \pi^0$ and $\bar{B}_s^0 \rightarrow K^0 \bar{K}^0$ and $\bar{B}_s^0 \rightarrow \eta \eta, \eta \eta'$ have not been observed yet. Some decays with small branching ratios expected from theoretical considerations, such as those decays, $\bar{B}^0 \rightarrow K^+ K^-$, $\bar{B}_s^0 \rightarrow \pi^+ \pi^-$, and $\bar{B}_s^0 \rightarrow \pi^0 \pi^0$ dominated by the annihilation contributions [11, 27] need further confirmation from data. Taking this positively, one can then use $SU(3)$ flavor symmetry framework to predict their branching ratios as further tests.

In this work, we will perform an updated global analysis for $B \rightarrow PP$ using the latest experimental data based on flavor symmetry. Without including η and η' in the final states, $SU(3)$ flavor symmetry is sufficient for the analysis. In order to include them also in the analysis, one needs to modify the analysis method. To this end we will enlarge the symmetry to $U(3)$ flavor symmetry, and also to take into account $\eta - \eta'$ mixing effect to study final states with P being η or η' . We find that $SU(3)$ flavor symmetry can explain data well without P being η or η' . The annihilation amplitudes are found to be small consistent with expectations. Several CP violating relations predicted by $SU(3)$ flavor symmetry are found in good agreement with data. Current available data can give constraints on the amplitudes which induce $P = \eta, \eta'$ decays in the framework of $U(3)$ flavor symmetry, and the $\eta - \eta'$ mixing angle θ can also be determined with $\theta = (-18.4 \pm 1.2)^\circ$ which is consistent with the value given by Particle Data Group from other fittings [1]. Several $B \rightarrow PP$ decay modes which have not been measured are predicted with branching ratios accessible at the LHCb. These decays can provide further tests for the framework of $SU(3)/U(3)$ flavor symmetry for B decays. In the following sections, we provide more details of our analysis.

II. $SU(3)$ DECAY AMPLITUDES FOR $B \rightarrow PP$

The quark level effective Hamiltonian responsible for charmless $B \rightarrow PP$ decays can be written as [28]

$$H_{eff}^q = \frac{4G_F}{\sqrt{2}} [V_{ub}V_{uq}^* (c_1 O_1 + c_2 O_2) - \sum_{i=3}^{11} (V_{ub}V_{uq}^* c_i^{uc} + V_{tb}V_{tq}^* c_i^{tc}) O_i], \quad (1)$$

with the superscript $q = d(s)$ for $\Delta S = 0 (-1)$ decay modes and V_{ij} the KM matrix elements. The coefficients $c_{1,2}$ and $c_i^{jk} = c_i^j - c_i^k$ are the Wilson Coefficients which have been evaluated by several groups [28] with $|c_{1,2}| \gg |c_i^{jk}|$. The operators O_i that consist of quarks and gluons can be written as

$$\begin{aligned} O_1 &= (\bar{q}_i u_j)_{V-A} (\bar{u}_i b_j)_{V-A}, & O_2 &= (\bar{q} u)_{V-A} (\bar{u} b)_{V-A}, \\ O_{3,5} &= (\bar{q} b)_{V-A} \sum_{q'} (\bar{q}' q')_{V \mp A}, & O_{4,6} &= (\bar{q}_i b_j)_{V-A} \sum_{q'} (\bar{q}'_j q'_i)_{V \mp A}, \\ O_{7,9} &= \frac{3}{2} (\bar{q} b)_{V-A} \sum_{q'} e_{q'} (\bar{q}' q')_{V \pm A}, & O_{8,10} &= \frac{3}{2} (\bar{q}_i b_j)_{V-A} \sum_{q'} e_{q'} (\bar{q}'_j q'_i)_{V \pm A}, \\ O_{11} &= \frac{g_s}{16\pi^2} \bar{q} \sigma_{\mu\nu} G^{\mu\nu} (1 + \gamma_5) b, & O_{12} &= \frac{Q_{be}}{16\pi^2} \bar{q} \sigma_{\mu\nu} F^{\mu\nu} (1 + \gamma_5) b. \end{aligned} \quad (2)$$

Under $SU(3)$ flavor symmetry, while the Lorentz-Dirac structure and color index are both omitted, $O_{1,2}$, $O_{3-6,11}$, and O_{7-10} transform as $\bar{3} + \bar{3}' + 6 + \bar{15}$, $\bar{3}$, and $\bar{3} + \bar{3}' + 6 + \bar{15}$, respectively [6–11]. As a result, H_{eff}^q can be decomposed as the matrices of $H(\bar{3})$, $H(6)$, and $H(\bar{15})$ with their non-zero entries to be [11]

$$\begin{aligned} H(\bar{3})^2 &= 1, \quad H(6)_1^{12} = H(6)_3^{23} = 1, \quad H(6)_1^{21} = H(6)_3^{32} = -1, \\ H(\bar{15})_1^{12} &= H(\bar{15})_1^{21} = 3, \quad H(\bar{15})_2^{22} = -2, \quad H(\bar{15})_3^{32} = H(\bar{15})_3^{23} = -1, \end{aligned} \quad (3)$$

for $\Delta S = 0$, and

$$\begin{aligned} H(\bar{3})^3 &= 1, \quad H(6)_1^{13} = H(6)_2^{32} = 1, \quad H(6)_1^{31} = H(6)_2^{23} = -1, \\ H(\bar{15})_1^{13} &= H(\bar{15})_1^{31} = 3, \quad H(\bar{15})_3^{33} = -2, \quad H(\bar{15})_2^{32} = H(\bar{15})_2^{23} = -1, \end{aligned} \quad (4)$$

for $\Delta S = -1$. Accordingly, the B mesons are presented as $B_i = (B_u, B_d, B_s) = (B^-, \bar{B}^0, \bar{B}_s^0)$, and for the final state P as the octet of $SU(3)$ representation M_j^i is given by

$$(M_j^i) = \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta_8}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta_8}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -2\frac{\eta_8}{\sqrt{6}} \end{pmatrix},$$

along with η_1 as the singlet of $SU(3)$ to be $(M_{\eta_1})_i^j = \delta_i^j \eta_1$. Note that $\bar{M} = M + M_{\eta_1}/\sqrt{3}$ form a nonet of $U(3)$. Consequently, without appealing to the dynamics of perturbative QCD, the $B \rightarrow PP$ decay amplitudes are given by

$$A(B \rightarrow PP) = \langle PP | H_{eff}^q | B \rangle = \frac{G_F}{\sqrt{2}} [V_{ub} V_{uq}^* T + V_{tb} V_{tq}^* P], \quad (5)$$

where the tree amplitude T for $B \rightarrow PP$ can be parametrized by $SU(3)$ invariant amplitudes. If one wants to include η_1 and η_8 into consideration, one may want to enlarge the analysis with $U(3)$ flavor symmetry. The $SU(3)/U(3)$ invariant amplitudes are given below ¹

$$\begin{aligned} T = & A_3^T B_i H(\bar{3})^i (\bar{M}_l^k \bar{M}_k^l) + C_3^T B_i \bar{M}_k^i \bar{M}_j^k H(\bar{3})^j \\ & + \tilde{A}_6^T B_i H(6)_k^{ij} \bar{M}_j^l \bar{M}_l^k + \tilde{C}_6^T B_i \bar{M}_j^i H(6)_l^{jk} \bar{M}_k^l \\ & + A_{\bar{15}}^T B_i H(\bar{15})_k^{ij} \bar{M}_j^l \bar{M}_l^k + C_{\bar{15}}^T B_i \bar{M}_j^i H(\bar{15})_l^{jk} \bar{M}_k^l \\ & + B_3^T B_i H(\bar{3})^i \bar{M}_j^j \bar{M}_k^k + \tilde{B}_6^T B_i H(6)_k^{ij} \bar{M}_j^k \bar{M}_l^l \\ & + B_{\bar{15}}^T B_i H(\bar{15})_k^{ij} \bar{M}_j^k \bar{M}_l^l + D_3^T B_i \bar{M}_j^i H(\bar{3})^j \bar{M}_l^l, \end{aligned} \quad (6)$$

with \tilde{C}_6 and \tilde{A}_6 rearranged to be $C_6 = \tilde{C}_6 - \tilde{A}_6$ [6–11]. Expanding the T expressions in Eq. (6), we obtain the tree amplitudes T in terms of the symmetry invariant amplitudes without η_8 and η_1 in the final states in Table I, while those with η_8 or/and η_1 in the final states are given in Table II. Note that the penguin amplitude P can be given by the replacement of the notation of T by P in the T amplitude, such that the hadronic parameters can be $C_{\bar{3},6,\bar{15}}^P$, $A_{\bar{3},\bar{15}}^P$, $B_{\bar{3},6,\bar{15}}^P$, and $D_{\bar{3}}^P$.

The dynamics of the interactions are all lumped into the invariant amplitudes, one cannot calculated the values for A_i , B_i , C_i , and D_i just from symmetry considerations, and therefore in our later analysis we will rely on experimental data to determine them. Note that $A_i^{T,P}$, $B_i^{T,P}$ are referred as annihilation amplitudes because the B mesons are first annihilated by the interaction Hamiltonian and two light mesons are then created and are expected to be smaller than C_i and D_i amplitudes.

Further simplification can be made because the operators for the tree and penguin contributions for the representations of 6 and $\bar{15}$, have the same structure, the differences for

¹ By treating η_1 as a $SU(3)$ singlet, we can form another T amplitude with $T = T_{\eta_8} + T_{\eta_1}$. Note that T_{η_8} can be given by using T in Eq. (6) where $\bar{M} = M + M_{\eta_1}/\sqrt{3}$ is replaced by $\bar{M} = M$, while T_{η_1} can be written as [18]

$$T_{\eta_1} = a^T B_i H(\bar{3})^i \eta_1 \eta_1 + b^T B_i M_j^i H(\bar{3})^j \eta_1 + c^T B_i H(6)_l^{ik} M_k^l \eta_1 + d^T B_i H(\bar{15})_l^{ik} M_k^l \eta_1.$$

The a^i , b^i , c^i , d^i and D^i , B^i amplitudes are related.

related amplitudes are caused by differences of the Wilson Coefficients (WC) in the Hamiltonian. Using WC obtained in Ref. [28], we use the numerical relations obtained in Ref. [18], $C_6^P(B_6^P) = -0.013C_6^T(B_6^T)$, and $C_{15}^P(A_{15}^P, B_{15}^P) + 0.015C_{15}^T(A_{15}^T, B_{15}^T)$, respectively. We comment that in finite order perturbative calculations the above relations are renormalization scheme and scale dependent. One should use a renormalization scheme consistently. We have checked with different renormalization schemes and find that numerically the changes are less than 15% for different schemes. In our later analysis, we will use the above relation. Moreover, since an overall phase can be removed without loss of generality, by setting C_3^P to be real, there can be totally 25 real independent parameters for $B \rightarrow PP$ in the SM with $SU(3)/U(3)$ flavor symmetry, given by

$$\begin{aligned} & C_3^P, \quad C_3^T e^{i\delta_3}, \quad C_6^T e^{i\delta_6}, \quad C_{15}^T e^{i\delta_{15}}, \quad A_3^T e^{i\delta_{A_3^T}}, \quad A_3^P e^{i\delta_{A_3^P}}, \quad A_{15}^T e^{i\delta_{A_{15}^T}}, \\ & B_3^T e^{i\delta_{B_3^T}}, \quad B_3^P e^{i\delta_{B_3^P}}, \quad B_6^T e^{i\delta_{B_6^T}}, \quad B_{15}^T e^{i\delta_{B_{15}^T}}, \quad D_3^T e^{i\delta_{D_3^T}}, \quad D_3^P e^{i\delta_{D_3^P}}. \end{aligned}$$

To obtain the amplitudes for B decays with at least one $\eta(\eta')$ in the final states, one also needs to consider $\eta_1 - \eta_8$ mixing,

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} \eta_8 \\ \eta_1 \end{pmatrix}, \quad (7)$$

where θ can be determined by fitting $B \rightarrow PP$ data.

III. NUMERICAL ANALYSIS AND DISCUSSIONS

In this section we carry out a global fit for $B \rightarrow PP$ using available experimental data to determine the $SU(3)/U(3)$ invariant amplitudes. In the numerical analysis we use Cabbibo-Kobayashi-Maskawa (CKM) parameters determined from other global analysis. We summarize the Wolfenstein parameters which determine CKM matrix elements in the following [1]

$$\lambda = 0.22543 \pm 0.00094, \quad A = 0.802 \pm 0.029, \quad \rho = 0.154 \pm 0.0124, \quad \eta = 0.363 \pm 0.0078.$$

For experimental inputs of the branching ratios and CP violating asymmetries, we use the data in Refs. [1, 2], while for $\mathcal{B}(B_d \rightarrow \pi^0 \eta)$ and $\mathcal{B}(B_s \rightarrow \eta' \eta')$ we use the newly observed ones from Refs. [29, 30], respectively.

To understand the significance of each type of amplitudes in explaining the data, we consider several different ways to carry out our numerical analysis. To see if indeed the

TABLE I. Decay amplitudes for $B \rightarrow PP$ without η_8 and η_1 .

$\Delta S = 0$	$\Delta S = -1$
$T_{\pi^-\pi^0}^{B_u}(d) = \frac{8}{\sqrt{2}}C_{15}^T,$	$T_{\pi^0K^-}^{B_u}(s) = \frac{1}{\sqrt{2}}(C_3^T - C_6^T + 3A_{15}^T + 7C_{15}^T),$
$T_{K^-K^0}^{B_u}(d) = C_3^T - C_6^T + 3A_{15}^T - C_{15}^T,$	$T_{\pi^-\bar{K}^0}^{B_u}(s) = C_3^T - C_6^T + 3A_{15}^T - C_{15}^T,$
$T_{\pi^+\pi^-}^{B_d}(d) = 2A_3^T + C_3^T + C_6^T + A_{15}^T + 3C_{15}^T,$	$T_{K^+\bar{K}^-}^{B_s}(s) = 2A_3^T + C_3^T + C_6^T + A_{15}^T + 3C_{15}^T,$
$T_{K^-K^+}^{B_d}(d) = 2(A_3^T + A_{15}^T),$	$T_{\pi^0\pi^0}^{B_s}(s) = \sqrt{2}(A_3^T + A_{15}^T),$
$T_{\pi^0\pi^0}^{B_d}(d) = \frac{1}{\sqrt{2}}(2A_3^T + C_3^T + C_6^T + A_{15}^T - 5C_{15}^T),$	$T_{K^0\bar{K}^0}^{B_s}(s) = 2A_3^T + C_3^T - C_6^T - 3A_{15}^T - C_{15}^T,$
$T_{\bar{K}^0K^0}^{B_d}(d) = 2A_3^T + C_3^T - C_6^T - 3A_{15}^T - C_{15}^T,$	$T_{\pi^+\pi^-}^{B_s}(s) = 2(A_3^T + A_{15}^T),$
$T_{K^0\pi^0}^{B_s}(d) = -\frac{1}{\sqrt{2}}(C_3^T + C_6^T - A_{15}^T - 5C_{15}^T),$	$T_{\pi^0\bar{K}^0}^{B_d}(s) = -\frac{1}{\sqrt{2}}(C_3^T + C_6^T - A_{15}^T - 5C_{15}^T),$
$T_{K^+\pi^-}^{B_s}(d) = C_3^T + C_6^T - A_{15}^T + 3C_{15}^T,$	$T_{\pi^+K^-}^{B_d}(s) = C_3^T + C_6^T - A_{15}^T + 3C_{15}^T.$

annihilation contributions are smaller than non-annihilation amplitudes, we analyze the data in two different ways: with or without annihilation contributions. The analysis with or without η and/or η' in the final states may also be significantly different because the mixing effect of $\eta - \eta'$ may complicate the situation. We therefore also carry out analysis according to whether or not include η and/or η' in the final states. In the case with η and/or η' in the final states, by fitting data, one may also obtain some information about the mixing angle θ . This may provide another way to determine the mixing angle.

Our results are presented for four different cases:

- 1). Analysis without annihilation contributions and without η and/or η' in the final states.
- 2). Analysis with annihilation contributions and without η and/or η' in the final states.
- 3). Analysis without annihilation contributions and with η and/or η' in the final states.
- 4). Analysis with annihilation contributions and with η and/or η' in the final states.

The values of the minimal χ^2 per degrees of freedom (d.o.f) for different cases from our fit are given by

$$\text{Case 1), } 1.65; \quad \text{Case 2), } 1.27; \quad \text{Case 3), } 1.71; \quad \text{Case 4), } 1.67. \quad (8)$$

TABLE II. Decay amplitudes for $B \rightarrow PP$ with at least one of the P being a η_8 or η_1 .

$\Delta S = 0$	$\Delta S = -1$
$T_{\pi^-\eta_8}^{B_u}(d) = \frac{2}{\sqrt{6}}(C_3^T - C_6^T + 3A_{15}^T + 3C_{15}^T),$	$T_{\eta_8 K^-}^{B_u}(s) = \frac{1}{\sqrt{6}}(-C_3^T + C_6^T - 3A_{15}^T + 9C_{15}^T),$
$T_{\pi^0\eta_8}^{B_d}(d) = \frac{1}{\sqrt{3}}(-C_3^T + C_6^T + 5A_{15}^T + C_{15}^T),$	$T_{\eta_8 \bar{K}^0}^{B_d}(s) = -\frac{1}{\sqrt{6}}(C_3^T + C_6^T - A_{15}^T - 5C_{15}^T),$
$T_{\eta_8\eta_8}^{B_d}(d) = \frac{1}{\sqrt{2}}(2A_3^T + \frac{1}{3}C_3^T - C_6^T - A_{15}^T + C_{15}^T),$	$T_{\pi^0\eta_8}^{B_s}(s) = \frac{2}{\sqrt{3}}(C_6^T + 2A_{15}^T - 2C_{15}^T),$
$T_{K^0\eta_8}^{B_s}(d) = -\frac{1}{\sqrt{6}}(C_3^T + C_6^T - A_{15}^T - 5C_{15}^T),$	$T_{\eta_8\eta_8}^{B_s}(s) = \sqrt{2}(A_3^T + \frac{2}{3}C_3^T - A_{15}^T - 2C_{15}^T),$
$T_{\pi^-\eta_1}^{B_u}(d) = \frac{1}{\sqrt{3}}(2C_3^T + C_6^T + 6A_{15}^T + 3C_{15}^T$ $+ 3B_6^T + 9B_{15}^T + 3D_3^T),$	$T_{K^-\eta_1}^{B_d}(s) = \frac{1}{\sqrt{3}}(2C_3^T + C_6^T + 6A_{15}^T + 3C_{15}^T$ $+ 3B_6^T + 9B_{15}^T + 3D_3^T),$
$T_{\pi^0\eta_1}^{B_d}(d) = \frac{-1}{\sqrt{6}}(2C_3^T + C_6^T - 10A_{15}^T - 5C_{15}^T$ $+ 3B_6^T - 15B_{15}^T + 3D_3^T),$	$T_{K^0\eta_1}^{B_d}(s) = \frac{1}{\sqrt{3}}(2C_3^T - C_6^T - 2A_{15}^T - C_{15}^T$ $- 3B_6^T - 3B_{15}^T + 3D_3^T),$
$T_{\eta_1\eta_8}^{B_d}(d) = \frac{1}{3\sqrt{2}}(2C_3^T - 3C_6^T + 6A_{15}^T + 3C_{15}^T$ $- 9B_6^T + 9B_{15}^T + 3D_3^T),$	$T_{\pi^0\eta_1}^{B_s}(s) = \frac{-2}{\sqrt{6}}(C_6^T - 4A_{15}^T - 2C_{15}^T$ $+ 3B_6^T - 6B_{15}^T),$
$T_{\eta_1\eta_1}^{B_d}(d) = \frac{\sqrt{2}}{3}(3A_3^T + C_3^T + 9B_3^T + 3D_3^T),$	$T_{\eta_1\eta_8}^{B_s}(s) = \frac{-\sqrt{2}}{3}(2C_3^T - 6A_{15}^T - 3C_{15}^T$ $- 9B_{15}^T + 3D_3^T),$
$T_{K^0\eta_1}^{B_s}(d) = \frac{1}{\sqrt{3}}(2C_3^T - C_6^T - 2A_{15}^T - C_{15}^T$ $- 3B_6^T - 3B_{15}^T + 3D_3^T),$	$T_{\eta_1\eta_1}^{B_s}(s) = \frac{\sqrt{2}}{3}(3A_3^T + C_3^T + 9B_3^T + 3D_3^T).$

Note that for each case, the minimal χ^2 is different because the available decay modes for data fitting for each case is different. The above minimal χ^2 per d.o.f indicate that all the four fits are reasonable ones.

The hadronic parameters determined for the four cases mentioned above are listed in Table III. After the hadronic parameters are determined, one can predict some of the not-yet-observed branching ratios and CP violating asymmetries. The results are given in Tables IV,V,VI,VII. In the following we comment on some features of our analysis.

As mentioned before, the annihilation contributions A_i are expected to be small compared with those of non-annihilation contributions C_i . Our fitting supports this expectation. The conclusions are drawn from comparing Case 1) with Case 2), and Case 3) with Case 4). Case 1) is an $SU(3)$ analysis neglecting annihilation contributions. A complete $SU(3)$ analysis would involve η_8 . However, due to $\eta - \eta'$ mixing, one cannot obtain a complete information when η_1 is not included. But if one restricts the analysis to only include

TABLE III. The best fit values and their 68% C.L. ranges for the hadronic parameters in the four cases.

	without η and η'		with η and η'	
	Case 1)	Case 2)	Case 3)	Case 4)
C_3^P	0.142 ± 0.001	0.141 ± 0.001	0.145 ± 0.002	0.142 ± 0.001
C_3^T	-0.188 ± 0.017	-0.198 ± 0.026	-0.197 ± 0.018	-0.211 ± 0.027
C_6^T	0.259 ± 0.021	0.257 ± 0.025	0.245 ± 0.016	0.255 ± 0.021
$C_{\overline{15}}^T$	-0.143 ± 0.004	-0.141 ± 0.004	-0.144 ± 0.004	-0.142 ± 0.004
$\delta_{\overline{3}}$	$(-121 \pm 5)^\circ$	$(-135 \pm 6)^\circ$	$(-124 \pm 5)^\circ$	$(-140 \pm 6)^\circ$
δ_6	$(50 \pm 4)^\circ$	$(54 \pm 6)^\circ$	$(51 \pm 4)^\circ$	$(56 \pm 6)^\circ$
$\delta_{\overline{15}}$	$(169 \pm 4)^\circ$	$(171 \pm 4)^\circ$	$(165 \pm 3)^\circ$	$(172 \pm 3)^\circ$
A_3^T	—	-0.034 ± 0.015	—	-0.039 ± 0.014
A_3^P	—	-0.013 ± 0.002	—	-0.013 ± 0.002
$A_{\overline{15}}^T$	—	-0.025 ± 0.012	—	-0.020 ± 0.012
$\delta_{A_3^T}$	—	$(-23 \pm 29)^\circ$	—	$(-16 \pm 25)^\circ$
$\delta_{A_3^P}$	—	$(-120 \pm 16)^\circ$	—	$(-123 \pm 16)^\circ$
$\delta_{A_{\overline{15}}^T}$	—	$(-30 \pm 26)^\circ$	—	$(-14 \pm 27)^\circ$
D_3^P	—	—	-0.077 ± 0.007	-0.073 ± 0.008
D_3^T	—	—	0.272 ± 0.036	0.275 ± 0.053
$\delta_{D_3^P}$	—	—	$(-55 \pm 9)^\circ$	$(-55 \pm 10)^\circ$
$\delta_{D_3^T}$	—	—	$(-90 \pm 9)^\circ$	$(-92 \pm 9)^\circ$
B_6^T	—	—	—	0.099 ± 0.094
$B_{\overline{15}}^T$	—	—	—	-0.038 ± 0.016
$\delta_{B_6^T}$	—	—	—	$(75 \pm 55)^\circ$
$\delta_{B_{\overline{15}}^T}$	—	—	—	$(78 \pm 48)^\circ$
θ	—	—	$(-18.4 \pm 1.2)^\circ$	$(-18.8 \pm 1.2)^\circ$
$\chi^2/d.o.f.$	1.65	1.27	1.71	1.67

pions and Kaons in the final state, the analysis should give a reasonable fit if the annihilation contributions are indeed small. This is indeed supported by the smallness of the branching ratios for those decays that only receive annihilation contributions, such as $B_d \rightarrow K^- K^+$,

$B_s \rightarrow \pi^+\pi^-$, and $B_s \rightarrow \pi^0\pi^0$. These modes only have branching ratios of order 10^{-7} . Analysis of Case 2) then helps to quantify the statement and obtain values for the relevant annihilation amplitudes. One can see that the annihilation amplitudes A_i are several times smaller than the non-annihilation amplitudes C_i . The comparison of Case 3) with Case 4) also supports this conclusion. From Table III, one can see that the current data still leave the amplitude D_i and B_i with large errors. We hope that when more data become available, the D_i and B_i amplitudes will have better accuracy and the expectation that annihilation contributions are smaller than non-annihilation contributions will be tested further in the sector involving η and η' in $B \rightarrow PP$ decays.

In case 3), there are 35 data points available with minimal $\chi^2/d.o.f$ of 1.71. The LHCb has measured many more decay modes compared with that could be achieved by using data from Belle and Babar detectors at B -factories only. In this case analysis with η and η' in the final states can be meaningfully carried out. One can even obtain information about the $\eta - \eta'$ mixing angle. The $\eta - \eta'$ mixing angle determined from Case 3) analysis gives $\theta = (-18.4 \pm 1.2)^\circ$. This is consistent with the value of $(-18 \pm 2)^\circ$ given by Particle Data Group [1].

Currently, the branching ratios and CP asymmetries for many decay modes with η and η' in the final states have not been observed, such as $\mathcal{B}(B_d \rightarrow \eta\eta, \eta\eta', \eta'\eta')$ and $\mathcal{B}(B_s \rightarrow \eta\eta, \eta\eta')$. Therefore, the theoretical predictions can be useful. For Case 3), the new parameters needed are D_i . The values for them are given in Table III. With the fitted D_i , we obtain $\mathcal{B}(B_u \rightarrow K^-\eta')$ and $\mathcal{B}(B_d \rightarrow \bar{K}^0\eta')$ to be $(75.0^{+2.3}_{-2.7}, 65.0^{+2.7}_{-2.5}) \times 10^{-6}$ which are consistent with data. We note that $\mathcal{B}(B_s \rightarrow \eta\eta')$ around 24×10^{-6} can be as large as the observed $\mathcal{B}(B_s \rightarrow \eta'\eta') = (33 \pm 11) \times 10^{-6}$, while $\mathcal{B}(B_d \rightarrow \eta\eta, \eta'\eta')$ of order 10^{-7} agree with the experimental upper bounds. When more data become available, this can be settled with high confidence.

In case 4), the parameters B_i with their phases, in principle, should be introduced implying 8 new parameters. We find that the determinations of $B_3^T e^{i\delta_{B_3^T}}$ and $B_3^P e^{i\delta_{B_3^P}}$ require at least 4 data points from $B_{d,s} \rightarrow \eta\eta, \eta\eta', \eta'\eta'$ decay modes, but only $\mathcal{B}(B_s \rightarrow \eta'\eta')$ is available. Present available data cannot determine $B_3^T e^{i\delta_{B_3^T}}$ and $B_3^P e^{i\delta_{B_3^P}}$. Since they are annihilation amplitudes which are expected to be small, we hence neglect their contributions for the practical fitting. Therefore, in this case we will have 21 parameters to fit 36 available data points. We obtain minimal $\chi^2/d.o.f$ to be 1.67 representing a reasonable fit. Again in this case, we can determine the $\eta - \eta'$ mixing angle θ with $\theta = (-18.8 \pm 1.2)^\circ$ represented

to be stable compared to that in case 3). The fitted B_i have larger uncertainties, such as $B_6^T = 0.099 \pm 0.094$. This is because that the data are not sufficient for the decays with η_1 , while A_i , B_i , C_i , and D_i are fitted together. When more data become available, the predictions made for this case can be tests; in particular, data will tell whether the omission of $B_3^T e^{i\delta_{B_3^T}}$ and $B_3^P e^{i\delta_{B_3^P}}$ for the fit is reasonable.

We now comment on a class of CP violating relations in the framework of $SU(3)$ flavor symmetry. This class of relations concern the rate difference among some B decays defined by

$$\Delta(B \rightarrow PP) = \Gamma(B \rightarrow PP) - \Gamma(\bar{B} \rightarrow \bar{P}\bar{P}), \quad (9)$$

which connects the branching ratio and the CP violating asymmetry with $\Delta(B_i \rightarrow PP) = \mathcal{A}_{CP}(B_i \rightarrow PP)\mathcal{B}(B_i \rightarrow PP)/\tau_{B_i}$ with τ_{B_i} the B_i lifetime.

The unique feature of the SM in the CKM matrix elements that $Im(V_{ub}V_{ud}^*V_{tb}^*V_{td}) = -Im(V_{ub}V_{us}^*V_{tb}^*V_{ts})$ can be used to relate the $\Delta S = 0$ and $\Delta S = -1$ decay modes with the same tree amplitude T and penguin amplitude P which can be read off from Table I. For instance, for $B_s \rightarrow K^+\pi^-$ and $B_d \rightarrow \pi^+K^-$, we obtain

$$\frac{\mathcal{A}_{CP}(B_d \rightarrow \pi^+K^-)}{\mathcal{A}_{CP}(B_s \rightarrow K^+\pi^-)} + \mathcal{R}(\Delta_{\pi^+K^-}^{B_d}/\Delta_{K^+\pi^-}^{B_s}) \frac{\mathcal{B}(B_s \rightarrow K^+\pi^-)/\tau_{B_s}}{\mathcal{B}(B_d \rightarrow \pi^+K^-)/\tau_{B_d}} = 0, \quad (10)$$

with $\mathcal{R}(\Delta_{\pi^+K^-}^{B_d}/\Delta_{K^+\pi^-}^{B_s}) = 1$.

If annihilation amplitudes are neglected, there are additional relations, for example

$$\frac{\mathcal{A}_{CP}(B_d \rightarrow \pi^+K^-)}{\mathcal{A}_{CP}(B_d \rightarrow \pi^+\pi^-)} + \mathcal{R}'(\Delta_{\pi^+K^-}^{B_d}/\Delta_{\pi^+\pi^-}^{B_d}) \frac{\mathcal{B}(B_d \rightarrow \pi^+\pi^-)}{\mathcal{B}(B_d \rightarrow \pi^+K^-)} \simeq 0, \quad (11)$$

with $\mathcal{R}'(\Delta_{\pi^+K^-}^{B_d}/\Delta_{\pi^+\pi^-}^{B_d}) \simeq 1$.

Deviation of \mathcal{R}_i away from 1 is a measure of $SU(3)$ flavor symmetry breaking. In Table VIII we list \mathcal{R}_i and \mathcal{R}'_i for some relations predicted with annihilation amplitudes and with annihilation amplitudes neglected, respectively. QCD based perturbation theory also predict similar values [26, 33, 36]. Note that experimentally, $\mathcal{R}_{data}(\Delta_{\pi^+K^-}^{B_d}/\Delta_{K^+\pi^-}^{B_s}) = 1.12 \pm 0.22$ and $\mathcal{R}'_{data}(\Delta_{\pi^+K^-}^{B_d}/\Delta_{\pi^+\pi^-}^{B_d}) \simeq 1.02 \pm 0.19$. The $SU(3)$ predictions are in good agreement with data. Since the relation with annihilation contributions neglected is also in good agreement with data, this also provides an evidence that annihilation contributions are indeed small. If $SU(3)$ is exact the fitted central value for \mathcal{R}_i should be equal to 1. The deviation in Table VIII is due to the fact that in calculating the values, we have used physics Kaon and pion

masses, branching ratios from fit and also experimental values for the lifetimes which slightly breaks $SU(3)$ flavor symmetry. Theoretically there are also several other pairs obeying the relations discussed (listed in Table VIII), at this moment there are large error bars to draw any conclusion. But once relevant quantities are measured, they will further test the theory.

In Table VII, we notice that several CP asymmetries are determined to be large. This is because accidental cancellations in the amplitudes for relevant decays (large final state interaction phase) and need to be tested. This may also reflect the fact that data are not sufficient to constrain the amplitudes with high precision and the “best” fits are are some very shallow local minimuns. More data are required to draw meaningful conclusions.

Finally, we make a comment on the recent theoretical study in Ref. [32] based on the diagrammatic $SU(3)$ flavor symmetry. Our fittings include the newly observed $\mathcal{B}(B_d \rightarrow \pi^0 \eta)$ and $\mathcal{B}(B_s \rightarrow \eta' \eta')$. Despite of the measured $\mathcal{B}(B_d \rightarrow \eta \eta') < 1.2 \times 10^{-6}$, we predict $\mathcal{B}(B_d \rightarrow \eta \eta')$ to be 2×10^{-6} similar to that in Ref. [32]. There is some tension between the fitted $\mathcal{B}(B_d \rightarrow \pi^0 \eta) = (0.91 \pm 0.03) \times 10^{-6}$ and the value of $(0.41 \pm 0.22) \times 10^{-6}$ from the data in comparison with $\mathcal{B}(B_d \rightarrow \pi^0 \eta) = (0.12 \pm 0.07) \times 10^{-6}$ [32]. Note that the predictions for $\mathcal{B}(B_d \rightarrow \pi^0 \eta)$ in the approaches of QCD factorization, pQCD, and SCET [33–35] are of order 10^{-8} . Future experiments can provide information to test these predictions.

IV. CONCLUSIONS

In this work we have performed an updated global analysis for $B \rightarrow PP$ using the latest experimental data based on flavor symmetry. Without including η and η' in the final states, $SU(3)$ flavor symmetry is sufficient for the analysis. In order to include P being η or η' in the analysis, we enlarged the symmetry to $U(3)$ flavor symmetry. In this case we also took into account $\eta - \eta'$ mixing effect. We found that $SU(3)$ flavor symmetry can explain data well without P being η or η' .

We have considered four different scenarios for data fitting to see how annihilation and also how inclusion of η and η' affect the results. The annihilation amplitudes were found to be small consistent with expectations. Current available data could give constraints on the amplitudes which induce $P = \eta, \eta'$ decays in the framework of $U(3)$ flavor symmetry. The $\eta - \eta'$ mixing angle θ could also be determined with $\theta = (-18.4 \pm 1.2)^\circ$ which is consistent with the value given by Particle Data Group from other fittings [1]. Several CP violating

relations predicted by $SU(3)$ flavor symmetry were found in good agreement with data. Although current data could not fix two annihilation amplitudes $B_3^{T,P} e^{i\delta_{B_3^{T,P}}}$, as they were expected to be small, we were able to predict several $B \rightarrow PP$ decay modes which have not been measured. These predicted branching ratios are accessible at the LHCb. We look forward to more data to come to test the framework of $SU(3)/U(3)$ flavor symmetry for B decays.

ACKNOWLEDGMENTS

The work was supported in part by MOE Academic Excellent Program (Grant No: 102R891505) and MOST of ROC, and in part by NSFC(Grant No:11175115 and 11575111) and Shanghai Science and Technology Commission (Grant No: 11DZ2260700) of PRC. We thank Tom Browder for bringing Ref.[37] to our attention.

Note Added: After submitting our paper to arXiv, we became aware that $B_s \rightarrow K^0 \bar{K}^0$ has been measured recently by Belle collaboration [37] with a branching ratio of $(19.6_{-5.1}^{+5.8}(stat.) \pm 1.0(sys.) \pm 2.0(N_{B_s^0 \bar{B}_s^0})) \times 10^{-6}$. The measured branching ratio in good agreement with our prediction.

-
- [1] K.A. Olive *et al.* [Particle Data Group Collaboration], Chin. Phys. C **38**, 090001 (2014).
 - [2] Y. Amhis *et al.*, arXiv:1412.7515 and online update at <http://www.slac.stanford.edu/xorg/hfag>.
 - [3] M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, Nucl. Phys. B **606**, 245 (2001); Nucl. Phys. B **675**, 333 (2003).
 - [4] Y.Y. Keum, H.n. Li and A.I. Sanda, Phys. Lett. B **504**, 6 (2001); Phys. Rev. D **63**, 054008 (2001).
 - [5] C.W. Bauer, S. Fleming, D. Pirjol and I.W. Stewart, Phys. Rev. D **63**, 114020 (2001); C.W. Bauer, D. Pirjol and I.W. Stewart, Phys. Rev. Lett. **87**, 201806 (2001).
 - [6] D. Zeppenfeld, Z. Phys. **C8**, 77(1981).

- [7] M.J. Savage and M.B. Wise, Phys. Rev. D **39**, 3346 (1989) [Phys. Rev. D **40**, 3127 (1989)].
- [8] L.L. Chau et al., Phys. Rev. **D43**, 2176 (1991).
- [9] M. Gronau *et al.*, Phys. Rev. **D50**, 4529 (1994); **D52**, 6356 (1995); *ibid*, 6374 (1995).
- [10] A.S. Dighe, M. Gronau and J.L. Rosner, Phys. Rev. Lett. **79**, 4333 (1997).
- [11] X.G. He, Eur. Phys. J. C **9**, 443 (1999).
- [12] N.G. Deshpande and X.G. He, Phys. Rev. Lett. **75**, 1703 (1995).
- [13] L.L. Chau, H.Y. Cheng, W.K. Sze, H. Yao and B. Tseng, Phys. Rev. D **43**, 2176 (1991) [Phys. Rev. D **58**, 019902 (1998)].
- [14] C.W. Chiang, M. Gronau, J.L. Rosner and D.A. Suprun, Phys. Rev. D **70**, 034020 (2004).
- [15] H.Y. Cheng and S. Oh, JHEP **1109**, 024 (2011).
- [16] W.S. Hou, H.n. Li, S. Mishima and M. Nagashima, Phys. Rev. Lett. **98**, 131801 (2007).
- [17] H.n. Li and S. Mishima, Phys. Rev. D **83**, 034023 (2011).
- [18] H.K. Fu, X.G. He and Y.K. Hsiao, Phys. Rev. D **69**, 074002 (2004).
- [19] X.G. He, G.N. Li and D. Xu, Phys. Rev. D **91**, 014029 (2015).
- [20] N.G. Deshpande and X.G. He, Phys. Rev. Lett. **75**, 3064 (1995).
- [21] X.G. He, Y.K. Hsiao, J.Q. Shi, Y.L. Wu and Y.F. Zhou, Phys. Rev. D **64**, 034002 (2001).
- [22] X.G. He, C.L. Hsueh and J.Q. Shi, Phys. Rev. Lett. **84**, 18 (2000).
- [23] G.N. Li, M. He and X.G. He, arXiv:1507.08252 [hep-ph].
- [24] M. He, X.G. He and G.N. Li, Phys. Rev. D **92**, 036010 (2015).
- [25] X.G. He and G.N. Li, Phys. Lett. B **750**, 82 (2015).
- [26] X.G. He, S.F. Li and H.H. Lin, JHEP **1308**, 065 (2013).
- [27] H.K. Fu, X.G. He, Y.K. Hsiao and J.Q. Shi, Chin. J. Phys. **41**, 601 (2003).
- [28] G.Buchalla, A.J. Buras, M.E. Lautenbacher, Rev. Mod. Phys. **68**, 1125(1996); A. Buras, M. Jamin, M. Lautenbacher, *ibid*, 75 (1993); M. Ciuchini et.al., Nucl. Phys. **B415**, 403 (1994); N.G. Deshpande and X.-G. He, Phys. Lett. **B336**, 417 (1994).
- [29] B. Pal *et al.* [BELLE Collaboration], Phys. Rev. D **92**, 011101 (2015).
- [30] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **115**, 051801 (2015).
- [31] H.K. Fu, X.G. He, Y.K. Hsiao and J.Q. Shi, Chin. J. Phys. **41**, 601 (2003); Nucl. Phys. Proc. Suppl. **115**, 279 (2003).
- [32] H.Y. Cheng, C.W. Chiang and A.L. Kuo, Phys. Rev. D **91**, 014011 (2015).
- [33] H.Y. Cheng and C.K. Chua, Phys. Rev. D **80**, 114026 (2009).

- [34] A. Ali, G. Kramer, Y. Li, C.D. Lu, Y.L. Shen, W. Wang and Y.M. Wang, Phys. Rev. D **76**, 074018 (2007).
- [35] A.R. Williamson and J. Zupan, Phys. Rev. D **74**, 014003 (2006); **74**, 03901(E) (2006).
- [36] H.Y. Cheng, C.K. Chua and A. Soni, Phys. Rev. D **72**, 014006 (2005).
- [37] B. Pal et al. (Belle Collaboration), arXiv:1512.02145[hep-ph].

TABLE IV. The central values and 68% C.L. allowed ranges for branching ratios (in units of 10^{-6}), where the superscript a denotes that the decay without C_i is not involved in the fitting.

Branching ratios	data	case1	case2	case3	case4
$B_u \rightarrow \pi^- \pi^0$	5.48 ± 0.35	$5.57^{+0.14}_{-0.13}$	$5.42^{+0.14}_{-0.13}$	$5.69^{+0.13}_{-0.13}$	$5.54^{+0.13}_{-0.12}$
$B_u \rightarrow K^- K^0$	1.32 ± 0.14	$1.34^{+0.04}_{-0.04}$	$1.34^{+0.08}_{-0.06}$	$1.20^{+0.04}_{-0.03}$	$1.18^{+0.07}_{-0.05}$
$B_d \rightarrow \pi^+ \pi^-$	5.10 ± 0.19	$5.20^{+0.14}_{-0.14}$	$5.12^{+0.22}_{-0.20}$	$5.22^{+0.14}_{-0.13}$	$5.13^{+0.23}_{-0.20}$
$B_d \rightarrow \pi^0 \pi^0$	1.17 ± 0.13	$1.05^{+0.04}_{-0.04}$	$1.15^{+0.06}_{-0.05}$	$1.06^{+0.04}_{-0.03}$	$1.17^{+0.05}_{-0.05}$
$B_d \rightarrow \bar{K}^0 K^0$	1.21 ± 0.16	$1.23^{+0.04}_{-0.03}$	$1.31^{+0.07}_{-0.05}$	$1.10^{+0.03}_{-0.03}$	$1.31^{+0.08}_{-0.06}$
$B_u \rightarrow \pi^- \bar{K}^0$	23.79 ± 0.75	$23.18^{+0.13}_{-0.13}$	$22.72^{+0.15}_{-0.14}$	$23.05^{+0.12}_{-0.12}$	$22.73^{+0.14}_{-0.14}$
$B_u \rightarrow \pi^0 K^-$	12.94 ± 0.52	$13.03^{+0.08}_{-0.08}$	$12.78^{+0.08}_{-0.08}$	$13.00^{+0.08}_{-0.08}$	$12.83^{+0.08}_{-0.08}$
$B_d \rightarrow \pi^+ K^-$	19.57 ± 0.53	$20.64^{+0.12}_{-0.12}$	$20.60^{+0.14}_{-0.13}$	$20.84^{+0.12}_{-0.12}$	$20.72^{+0.13}_{-0.12}$
$B_d \rightarrow \pi^0 \bar{K}^0$	9.93 ± 0.49	$9.20^{+0.06}_{-0.06}$	$9.15^{+0.06}_{-0.06}$	$9.28^{+0.06}_{-0.06}$	$9.20^{+0.06}_{-0.06}$
$B_d \rightarrow K^+ K^-$	0.13 ± 0.05	— ^a	$0.14^{+0.03}_{-0.02}$	— ^a	$0.14^{+0.03}_{-0.02}$
$B_s \rightarrow K^+ \pi^-$	5.5 ± 0.5	$5.0^{+0.1}_{-0.1}$	$5.57^{+0.19}_{-0.19}$	$5.01^{+0.13}_{-0.13}$	$5.61^{+0.20}_{-0.17}$
$B_s \rightarrow K^0 \pi^0$	—	$2.02^{+0.08}_{-0.07}$	$1.59^{+0.08}_{-0.07}$	$2.04^{+0.07}_{-0.07}$	$1.64^{+0.08}_{-0.06}$
$B_s \rightarrow K^+ K^-$	24.8 ± 1.7	$19.8^{+0.1}_{-0.1}$	$24.5^{+0.6}_{-0.6}$	$20.0^{+0.1}_{-0.1}$	$24.5^{+0.6}_{-0.6}$
$B_s \rightarrow K^0 \bar{K}^0$	< 66	$20.5^{+0.1}_{-0.1}$	$22.9^{+0.3}_{-0.3}$	$20.4^{+0.1}_{-0.1}$	$22.4^{+0.4}_{-0.3}$
$B_s \rightarrow \pi^+ \pi^-$	0.76 ± 0.19	— ^a	$0.72^{+0.06}_{-0.05}$	— ^a	$0.71^{+0.06}_{-0.05}$
$B_s \rightarrow \pi^0 \pi^0$	< 210	— ^a	$0.18^{+0.01}_{-0.01}$	— ^a	$0.18^{+0.01}_{-0.01}$

TABLE V. The central values and 68% C.L. allowed ranges for CP asymmetries (in units of 10^{-2}).

CP asymmetries	data	case1	case2	case3	case4
$B_u \rightarrow \pi^- \pi^0$	2.6 ± 3.9	0 ± 0	0 ± 0	0 ± 0	0 ± 0
$B_u \rightarrow K^- K^0$	-8.7 ± 10.0	$-2.8^{+4.0}_{-4.0}$	$-3.8^{+7.4}_{-6.8}$	$-5.5^{+3.8}_{-3.8}$	$-7.7^{+8.6}_{-7.2}$
$B_d \rightarrow \pi^+ \pi^-$	31 ± 5	$31.1^{+1.1}_{-1.1}$	$30.2^{+2.2}_{-2.4}$	$31.1^{+1.1}_{-1.1}$	$29.7^{+2.0}_{-2.1}$
$B_d \rightarrow \pi^0 \pi^0$	43.0 ± 24.0	$57.2^{+1.2}_{-1.3}$	$64.0^{+1.8}_{-1.9}$	$56.1^{+1.2}_{-1.2}$	$63.3^{+1.7}_{-1.8}$
$B_d \rightarrow \bar{K}^0 K^0$	-60.0 ± 70.0	$-2.8^{+4.0}_{-4.0}$	$-17.8^{+9.7}_{-8.6}$	$-5.5^{+4.0}_{-3.8}$	$-18.0^{+9.2}_{-8.1}$
$B_u \rightarrow \pi^- \bar{K}^0$	-1.7 ± 1.6	$0.17^{+0.24}_{-0.24}$	$0.23^{+0.64}_{-0.47}$	$0.30^{+0.22}_{-0.22}$	$0.42^{+0.39}_{-0.48}$
$B_u \rightarrow \pi^0 K^-$	4.0 ± 2.1	$5.8^{+0.5}_{-0.5}$	$4.2^{+0.4}_{-0.7}$	$5.8^{+0.4}_{-0.4}$	$4.7^{+0.6}_{-0.6}$
$B_d \rightarrow \pi^+ K^-$	-8.2 ± 0.6	$-7.8^{+0.3}_{-0.3}$	$-8.1^{+0.4}_{-0.4}$	$-7.9^{+0.3}_{-0.3}$	$-8.0^{+0.4}_{-0.4}$
$B_d \rightarrow \pi^0 \bar{K}^0$	0 ± 13	$-13.3^{+0.5}_{-0.5}$	$-11.3^{+0.5}_{-0.5}$	$-13.2^{+0.4}_{-0.4}$	$-11.6^{+0.5}_{-0.5}$
$B_d \rightarrow K^+ K^-$	—	— ^a	$82.8^{+4.4}_{-6.0}$	— ^a	$83.6^{+4.4}_{-6.2}$
$B_s \rightarrow K^+ \pi^-$	26.0 ± 4.0	$31.1^{+1.1}_{-1.2}$	$28.1^{+1.4}_{-1.3}$	$31.2^{+1.0}_{-0.1}$	$28.0^{+1.2}_{-1.1}$
$B_s \rightarrow K^0 \pi^0$	—	$57.2^{+1.2}_{-1.3}$	$61.4^{+1.8}_{-2.1}$	$55.9^{+1.2}_{-1.2}$	$60.6^{+1.6}_{-1.9}$
$B_s \rightarrow K^+ K^-$	-14 ± 11	$-8.0^{+0.3}_{-0.4}$	$-5.6^{+0.5}_{-0.5}$	$-8.0^{+0.3}_{-0.3}$	$-5.6^{+0.4}_{-0.5}$
$B_s \rightarrow K^0 \bar{K}^0$	—	$0.17^{+0.24}_{-0.23}$	$12.1^{+1.2}_{-1.3}$	$0.27^{+0.21}_{-0.22}$	$10.4^{+1.3}_{-1.4}$
$B_s \rightarrow \pi^+ \pi^-$	—	— ^a	$-16.1^{+1.9}_{-1.6}$	— ^a	$-16.2^{+2.1}_{-2.1}$
$B_s \rightarrow \pi^0 \pi^0$	< 210	— ^a	$-16.1^{+1.9}_{-1.9}$	— ^a	$-16.2^{+2.0}_{-2.0}$

TABLE VI. The central values and their 68% C.L. allowed ranges for branching ratios (in units of 10^{-6}) with at least one of the final mesons to be a η or η' .

Branching ratios	data	case3	case4
$B_u \rightarrow \pi^- \eta$	4.02 ± 0.27	$3.77^{+0.12}_{-0.11}$	$3.73^{+1.50}_{-0.45}$
$B_u \rightarrow \pi^- \eta'$	2.7 ± 0.5	$3.33^{+0.19}_{-0.16}$	$3.23^{+8.81}_{-0.92}$
$B_d \rightarrow \pi^0 \eta$	0.41 ± 0.22	$0.91^{+0.03}_{-0.03}$	$0.77^{+0.61}_{-0.02}$
$B_d \rightarrow \pi^0 \eta'$	1.2 ± 0.4	$1.06^{+0.06}_{-0.05}$	$1.23^{+4.21}_{-0.11}$
$B_u \rightarrow K^- \eta$	2.36 ± 0.22	$2.16^{+0.22}_{-0.17}$	$2.19^{+0.37}_{-0.24}$
$B_u \rightarrow K^- \eta'$	71.1 ± 2.6	$75.0^{+2.3}_{-2.7}$	$71.1^{+4.7}_{-3.6}$
$B_d \rightarrow \bar{K}^0 \eta$	1.23 ± 0.27	$1.63^{+0.19}_{-0.15}$	$1.54^{+0.28}_{-0.17}$
$B_d \rightarrow \bar{K}^0 \eta'$	66.1 ± 3.1	$65.0^{+2.7}_{-2.5}$	$64.5^{+4.2}_{-3.4}$
$B_d \rightarrow \eta \eta$	< 1.0	$0.33^{+0.02}_{-0.01}$	$0.55^{+0.84}_{-0.11}$
$B_d \rightarrow \eta \eta'$	< 1.2	$1.91^{+0.10}_{-0.10}$	$3.33^{+10.06}_{-0.66}$
$B_d \rightarrow \eta' \eta'$	< 1.7	$0.41^{+0.03}_{-0.02}$	$0.28^{+0.92}_{-0.02}$
$B_s \rightarrow K \eta$	—	$0.99^{+0.04}_{-0.04}$	$1.12^{+1.84}_{-0.29}$
$B_s \rightarrow K \eta'$	—	$3.52^{+0.16}_{-0.14}$	$4.29^{+10.29}_{-0.48}$
$B_s \rightarrow \pi^0 \eta$	< 1000	$0.048^{+0.002}_{-0.002}$	$0.037^{+0.13}_{-0.01}$
$B_s \rightarrow \pi^0 \eta'$	—	$0.085^{+0.003}_{-0.003}$	$0.25^{+1.24}_{-0.06}$
$B_s \rightarrow \eta \eta$	< 1500	$2.81^{+0.12}_{-0.11}$	$3.29^{+0.13}_{-0.06}$
$B_s \rightarrow \eta \eta'$	—	$23.70^{+0.65}_{-0.54}$	$21.99^{+0.58}_{-0.13}$
$B_s \rightarrow \eta' \eta'$	33.1 ± 10.4	$21.30^{+1.10}_{-0.90}$	$20.42^{+1.15}_{-1.00}$

TABLE VII. The central values and their 68% C.L. allowed ranges for CP asymmetries (in units of 10^{-2}) with at least one of the final mesons to be a η or η' .

CP asymmetries	data	case3	case4
$B_u \rightarrow \pi^- \eta$	-14 ± 5	$-14.6^{+2.8}_{-2.7}$	$-12.3^{+28.5}_{-20.9}$
$B_u \rightarrow \pi^- \eta'$	6 ± 15	$8.9^{+5.9}_{-6.3}$	$5.6^{+22.8}_{-23.4}$
$B_d \rightarrow \pi^0 \eta$	—	$-26.8^{+4.2}_{-3.9}$	$-0.4^{+30.4}_{-26.7}$
$B_d \rightarrow \pi^0 \eta'$	—	$-48.5^{+7.6}_{-6.5}$	$83.3^{+5.2}_{-57.6}$
$B_u \rightarrow K^- \eta$	-37 ± 8	$-30.9^{+2.3}_{-2.4}$	$-31.1^{+13.3}_{-9.9}$
$B_u \rightarrow K^- \eta'$	1.3 ± 1.7	$0.5^{+0.3}_{-0.3}$	$0.8^{+6.8}_{-7.5}$
$B_d \rightarrow \bar{K}^0 \eta$	—	$3.2^{+1.8}_{-2.2}$	$8.7^{+16.8}_{-12.2}$
$B_d \rightarrow \bar{K}^0 \eta'$	—	$4.3^{+0.3}_{-0.3}$	$34.8^{+7.4}_{-6.9}$
$B_d \rightarrow \eta \eta$	—	$-86.6^{+2.0}_{-1.6}$	$-42.1^{+53.1}_{-2.6}$
$B_d \rightarrow \eta \eta'$	—	$-68.8^{+5.4}_{-4.3}$	$-27.9^{+51.9}_{-6.7}$
$B_d \rightarrow \eta' \eta'$	—	$-62.7^{+6.4}_{-5.5}$	$-87.9^{+56.5}_{-10.8}$
$B_s \rightarrow K \eta$	—	$-5.5^{+3.4}_{-3.4}$	$-11.5^{+28.8}_{-13.4}$
$B_s \rightarrow K \eta'$	—	$-79.7^{+4.1}_{-3.1}$	$-93.0^{+62.6}_{-2.1}$
$B_s \rightarrow \pi^0 \eta$	—	$98.1^{+0.4}_{-0.7}$	$83.3^{+4.8}_{-57.3}$
$B_s \rightarrow \pi^0 \eta'$	—	$98.1^{+0.4}_{-0.7}$	$64.7^{+10.0}_{-35.4}$
$B_s \rightarrow \eta \eta$	—	$-13.5^{+0.4}_{-0.4}$	$6.0^{+2.1}_{-3.2}$
$B_s \rightarrow \eta \eta'$	—	$-3.1^{+0.3}_{-0.4}$	$-1.3^{+2.5}_{-1.3}$
$B_s \rightarrow \eta' \eta'$	—	$4.5^{+0.4}_{-0.4}$	$4.8^{+4.5}_{-3.7}$

TABLE VIII. $\mathcal{R}_i^{(\prime)}$ to test the $SU(3)$ flavor symmetry. The fitted numbers in the parentheses are for case 1) and case 2), respectively.

modes	\mathcal{R}_{data}	$\mathcal{R}_{fit}^{(\prime)}$
$\mathcal{R}(\Delta_{\pi^+ K^-}^{B_d} / \Delta_{K^+ \pi^-}^{B_s})$	1.12 ± 0.22	$(1.03 \pm 0.06, 1.06 \pm 0.08)$
$\mathcal{R}(\Delta_{K^+ K^-}^{B_s} / \Delta_{\pi^+ \pi^-}^{B_d})$	2.20 ± 1.77	$(0.98 \pm 0.06, 0.89 \pm 0.12)$
$\mathcal{R}(\Delta_{\pi^- \bar{K}^0}^{B_u} / \Delta_{K^- K^0}^{B_u})$	-3.52 ± 5.25	$(1.05 \pm 2.07, 1.02 \pm 3.48)$
$\mathcal{R}(\Delta_{\pi^0 \bar{K}^0}^{B_d} / \Delta_{K^0 \pi^0}^{B_s})$	—	$(1.06 \pm 0.06, 1.06 \pm 0.08)$
$\mathcal{R}(\Delta_{\pi^+ \pi^-}^{B_s} / \Delta_{K^- K^+}^{B_d})$	—	$(—, 1.00 \pm 0.27)$
$\mathcal{R}(\Delta_{\pi^0 \pi^0}^{B_s} / \Delta_{K^- K^+}^{B_d})$	—	$(—, 1.00 \pm 0.02)$
$\mathcal{R}'(\Delta_{\pi^+ K^-}^{B_d} / \Delta_{\pi^+ \pi^-}^{B_d})$	1.02 ± 0.19	$(0.99 \pm 0.06, 1.07 \pm 0.11)$
$\mathcal{R}'(\Delta_{\pi^0 \bar{K}^0}^{B_d} / \Delta_{\pi^0 \pi^0}^{B_d})$	0.00 ± 1.28	$(1.02 \pm 0.06, 0.70 \pm 0.05)$
$\mathcal{R}'(\Delta_{K^+ K^-}^{B_s} / \Delta_{K^+ \pi^-}^{B_s})$	2.42 ± 1.96	$(1.01 \pm 0.06, 0.88 \pm 0.10)$
$\mathcal{R}'(\Delta_{\pi^- \bar{K}^0}^{B_u} / \Delta_{\bar{K}^0 K^0}^{B_d})$	-0.56 ± 0.83	$(1.14 \pm 2.28, 0.22 \pm 0.64)$